

Lipid-analogous phosphoric triesters


## Description

5 The present invention relates to novel phosphoric triesters which comprise apolar lipid structures.

These triesters can be used in particular as liposome constituents.

10

One aspect of the invention is a compound of the formula (I)



15 in which R<sup>1</sup> is a residue selected from cholesterol, diacylglycerols, dialkylglycerols, acylalkylglycerols, ceramides, primary or secondary alcohols having 12 to 24 C atoms or acylglycerobenzyl ethers, R<sup>2</sup> is a residue selected from ethanolamine, N-methylethanolamine,  
20 propanolamine, choline, glycerol, oligoglycerols, glycoglycerols or serine, each of which may optionally comprise protective groups, and R<sup>3</sup> is a radical selected from C<sub>1</sub>-C<sub>8</sub>-alkyl or C<sub>1</sub>-C<sub>8</sub>-alkenyl or has the meaning of R<sup>2</sup>.

25


The compounds of the invention are lipid-analogous phosphoric triesters, in particular phosphoric triesters having multiple hydroxyl groups per phosphorus atom, in particular at least two hydroxyl  
30 groups per phosphorus atom, more preferably at least three hydroxyl groups per phosphorus atom and even more preferably at least four hydroxyl groups per phosphorus atom.

35 The phosphoric triesters of the invention comprise a residue R<sup>1</sup> which comprises an apolar lipid structure.

Suitable structures for R<sup>1</sup> are in particular cholesterol residues, so that preferred phosphoric triesters are cholesteryl compounds.

- 5 A further preferred residue for R<sup>1</sup> is diacylglycerol, where the acyl groups each comprise independently preferably 12 to 28, in particular 13 to 27 and more preferably 14 to 26 carbon atoms. The acyl radicals may be saturated or mono- or poly-, in particular di- or  
10 triunsaturated radicals. Acylglycerols which comprise unsaturated fatty acid residues such as, for example, residues of oleic acid, linoleic acid or linolenic acid are particularly preferred.
- 15 The residue R<sup>1</sup> may additionally according to the invention be a dialkylglycerol residue, wherein the alkyl radicals each independently of one another preferably have 1 to 28, in particular 12 to 26, even more preferably 14 to 24 carbon atoms. The alkyl  
20 radicals in the dialkylglycerol residue may be saturated or mono- or poly-, in particular di- or triunsaturated. Preferred radicals are (Z)-9-octadecenyl-, (Z.Z.)-9.12-octadecanedieryl-, (Z.Z.Z)-9.12.15-octadecanetrieryl-, and lipophilic basic  
25 structures with a pharmaceutical effect, such as 1-octadecyl-2-methyl-sn-glycerol.

R<sup>1</sup> may further be a ceramide residue. Ceramides are endogenous lipophilic amides which are to be found in  
30 particular bound in the cerebral matter and in the myelin of the CNS and have the general formula (IV)



where R<sup>4</sup> is a long-chain fatty acid residue, in  
35 particular a fatty acid residue having 12 to 28 C atoms, R<sup>5</sup> is a long-chain alkyl radical, in particular

an alkyl radical having 12 to 28 C atoms and  $R^6$  is H.

$R^1$  may further be a residue of a primary or secondary alcohol having 12 to 24 carbon atoms, in particular 13 to 22 carbon atoms, where the alcohols may be saturated or mono- or polyunsaturated.

$R^1$  may further be an acylglycerobenzyl ether residue, it being possible to employ such compounds in particular as starting materials for synthesizing lysophospholipids.

The residue  $R^1$  may be present in the compounds of the invention in enantiopure form or as racemic mixture.

$R^2$  can be all the residues occurring in natural phospholipids and sphingomyelins.  $R^2$  is in particular an ethanolamine residue, an N-methylethanolamine residue or a propanolamine residue, where the residues are provided where appropriate with suitable protective groups, for example BOC.  $R^2$  may further be a choline residue.  $R^2$  is preferably  $-\text{CH}_2-\text{CH}_2-\text{N}^+(\text{CH}_3)_3$ .  $R^2$  may further be a glycerol residue ( $-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_2(\text{OH})$ ) and be an oligoglycerol, in particular a di- or triglycerol residue. Further suitable  $R^2$  residues are glycoglycerols, and serine residues. The glycerol and serine residues may also where appropriate be provided with suitable protective groups.

$R^3$  is a radical selected from  $\text{C}_1$ - $\text{C}_8$ -alkyl or  $\text{C}_1$ - $\text{C}_8$ -alkenyl or may have one of the meanings indicated above for  $R^2$ . If  $R^3$  has one of the meanings indicated for  $R^2$ , it is possible to form highly biologically active structures which, as novel cationic lipids, have great importance. Such cationic lipids can be employed for example for gene transfection.


$R^3$  may, however, also have only a temporary character, that is to say assume the function of a protective

group which is detached again later. In this case, R<sup>3</sup> is preferably methyl, ethyl, allyl or propyl.

The compounds of the invention are very stable between  
5 pH 3 and pH 8 and can be used in particular as liposome constituents.


The invention further relates to a process for  
10 preparing a compound according to formula (I), which is characterized in that a compound of the formula (II) is esterified with a compound of the formula (III) HO-R<sup>3</sup>.

The compounds are derived from phospholipids and are produced for example from



15


by esterification of the phosphoric diester with glycerol:




20

Cholesterol derivatives can also be obtained

correspondingly



- These compounds can be prepared in various embodiments.
- 5 They may also comprise oligoglycerols, e.g. for example glyceroglycerol, diglyceroglycerol or triglyceroglycerol in place of glycerol. Diagrammatically, for example, cholesterol-phospho-monoglycerol-triglyceroglycerol has the following structural formula:



10

The compounds of the invention are particularly suitable for preparing liposomes and as liposome constituents. They confer particular properties on


15 liposomes, e.g. long circulation times in the blood, targeted enrichment in the liver or else almost exclusive uptake in the spleen. It is also possible with the aid of the phosphoric triesters of the invention to form liposomes with novel properties,

20 which have high serum stability, have long circulation times and accumulate exclusively in the spleen. Long

circulation times are, however, also particularly important because the structures then do not, like known liposomes, accumulate in the liver but may hit other targets such as, for example, the spleen or, particularly importantly, be taken up by tumor cells. The compounds of the invention can therefore also be employed for the treatment of cancers.

The invention further relates to a novel synthetic route using the phosphoric triesters of the invention as intermediate. A particular advantage of the synthetic route of the invention is that the reaction direction (a) used in earlier syntheses is avoided, and according to (b) the important compound 1,2-dioleoyl-sn-glycero-3-phosphoglycerol or corresponding compounds are liberated under neutral conditions:

$R_1 = 1,2\text{-dioleoyl-sn-glycerol}$



Central intermediate


20

A substantial advantage of the novel management of the synthesis is the possibility of advancing apolar intermediates as far as possible in the synthesis, so that polar structures are introduced only at the end of the process. This is illustrated below in an example. Cardiolipins and analogous compounds are complicated structures which can be obtained only with great difficulty by synthesis in kg quantities. However, with the aid of our novel synthetic strategy, this is easily

25

possible.


Cardiolipin



5

The synthesis is described for the example of R = palmitic acid. The starting material is 1,2-dipalmitoyl-sn-glycerol which is converted with phosphorus oxychloride in THF with triethylamine as base in the usual manner into 1,2-dipalmitoyl-sn-glycero-3-phosphoric dichloride:


10



Building block I

15


The simple route for synthesizing cardiolipin, direct reaction with 2-benzylglycerol, unfortunately leads to predominant formation of the corresponding phospholane and is not practicable:



It is therefore necessary to use building block II, a protected glycerol derivative:

5


## Building block II



Linkage of building block I with building block II in  
10 THF with triethylamine then leads to building block  
III:

### Building block III


15      Building block I      ↓      1) Building block II  
  (triethylamine in THF)  
  2) Methanolysis  
  ↓      3) H<sup>(+)</sup>/H<sub>2</sub>O in THF



20

Building block III can then be reacted with building block I in the usual way to give the direct precursor of cardiolipin, which is then converted by





5 methanolysis into the dimethyl ester. The hydroxyl group on the middle glycerol is then liberated by catalytic hydrogenolysis. The methyl groups are removed by LiBr at neutral pH - the final product is cardiolipin.

10 The description is illustrated further by the following examples.


#### Example 1

- 1) Cholesteryl-phospho-diglycerol  
C<sub>33</sub>H<sub>59</sub>O<sub>8</sub>P (MW 614.801)
- 2) Cholesteryl-phospho-glycerol-glyceroglycerol  
C<sub>36</sub>H<sub>65</sub>O<sub>10</sub>P (MW 688.880)
- 3) Cholesteryl-phospho-di-glycoglycerol  
C<sub>37</sub>H<sub>67</sub>O<sub>10</sub>P (MW 702.907)
- 4) 1.2-Dimyristoyl-sn-glycero-3-phospho-diglycerol  
C<sub>37</sub>H<sub>73</sub>O<sub>12</sub>P (MW 740.953)
- 5) 1.2-Dipalmitoyl-sn-glycero-3-phospho-diglycerol  
C<sub>41</sub>H<sub>81</sub>O<sub>12</sub>P (MW 797.061)

- 6) 1.2-Distearoyl-sn-glycero-3-phospho-diglycerol  
C<sub>45</sub>H<sub>89</sub>O<sub>12</sub>P (MW 853.169)
- 7) 1.2-Dioleoyl-sn-glycero-3-phospho-diglycerol  
C<sub>45</sub>H<sub>85</sub>O<sub>12</sub>P (MW 849.137)
- 8) 1.2-Dioleoyl-sn-glycero-3-phospho-di-glycoglycerol  
C<sub>49</sub>H<sub>93</sub>O<sub>14</sub>P (MW 937.243)
- 9) 1.2-Dioleoyl-sn-glycero-3-phospho-di-glyceroglycerol  
C<sub>51</sub>H<sub>97</sub>O<sub>16</sub>P (MW 997.295)

Example 2


- 1) R<sub>1</sub>: 1.2.-dimyristoyl-glycerol
- 5 R<sub>2</sub>: choline  
R<sub>3</sub>: glycerol



and corresponding structures with 1.2.dioleoylglycerol.

10

- 2) R<sub>1</sub>: 1.2-dioleoylglycerol  
R<sub>2</sub>: glycerol  
R<sub>3</sub>: methyl



### Example 3

Liposomes of the composition

5

	<u>Molar ratio</u>
1.2-Distearoyl-sn-glycero-3-phosphocholine	40%
Cholesterol	30%
Cholesterol-phospho-diglycerol	20%
Cholesterol-phospho-glycerol, Na <sup>(+)</sup> salt	<u>10%</u>
	100%

accumulate predominantly in the spleen, whereas liposomes of the usual composition e.g.

1.2-Distearoyl-sn-glycero-3-phosphocholine	50%
Cholesterol	40%
1.2-Distearoyl-sn-glycero-3-phosphoglycerol, Na <sup>(+)</sup> salt	<u>10%</u>
	100%

10

accumulate mainly in the liver.